

Far Eastern Entomologist

Дальневосточный энтомолог

Journal published by Far East Branch
of the Russian Entomological Society
and Laboratory of Entomology,
Institute of Biology and Soil Science,
Vladivostok

Number 191: 1-7

ISSN 1026-051X

December 2008

NEW RECORDS OF VELVET ANTS (HYMENOPTERA: MUTILIDAE) FROM THE WEST OF IRAN

A.S. Lelej¹⁾, B. Gharali²⁾, H. Lotfalizadeh³⁾

1) Institute of Biology and Soil Science, Far Eastern Branch of Russian Academy of Sciences, Vladivostok-22, 690022, Russia. E-mail: lelej@biosoil.ru

2) Department of Plant Protection, Ghazvin Research Center for Agriculture and Natural Resources, Shahid Beheshti Blvd., No. 118, P.O. Box: 34185-618, Ghazvin, Iran. E-mail: bgharaei@mail.com

3) Department of Insect taxonomy, Iranian Plant Protection Research Institute, P.O. Box: 19395-1454, Tehran, Iran. E-mail: hlotfalizadeh2001@yahoo.com

Eight species are recorded from the west of Iran. *Smicromyrme* (*Erimyrme*) *zagrosensis* Lelej **sp. n.** (Ilam Province) is described and figured. Four species are newly recorded from Iran.

KEY WORDS: Velvet ants, Mutilidae, fauna, taxonomy, new species, Iran.

А. С. Лелей¹⁾, Б. Гарали²⁾, Х. Лотфализаде³⁾. Новые данные об ос-немках из Западного Ирана // Дальневосточный энтомолог. 2008. N 191. С. 1-7.

Из Западного Ирана приведены 8 видов ос-немок. Описывается *Smicromyrme* (*Erimyrme*) *zagrosensis* Lelej **sp. n.** из провинции Илам. 4 вида впервые указываются для фауны Ирана.

1) Биолого-почвенный институт ДВО РАН, Владивосток-22, 690022, Россия.

2) Отдел защиты растений, Исследовательский центр сельского хозяйства и природных ресурсов, Газвин, 34185-618, Иран.

3) Отдел таксономии насекомых, Иранский институт защиты растений, Тегеран, 19395-1454, Иран.

INTRODUCTION

Mutillidae is a large family of solitary wasps which parasitoids on the enclosed immature of other insects (Brothers, 1989). Adults are covered with dense velvety hairs and females sting severely and painfully when they are disturbed. Velvet ants are often conspicuous and brightly colored black and scarlet, or black with orange or whitish hairs. In most species, the segments of mesosoma are fused together so that they are immobile and boxlike (Brothers & Finnamore, 1993). In the field, they are easily recognized by bright patterns on the abdomen and dense hairs and filiform antennae. Mutillidae currently includes 208 genera and about 4200 described species (Lelej & Brothers, 2008; Lelej, 2007). The estimated number of species is much higher, numbering approximately 6000. They are distributed in all biogeographical regions, their greatest diversity occurs in the tropical and subtropical regions of the world. Some species of velvet ants are nocturnal, but many are active during the day.

Fauna of these wasps has been reviewed in the Palaearctic rather well. The catalogue of Palaearctic Mutillidae (523 species in 54 genera) with the key to the genera has been published recently (Lelej, 2002). Sixty-seven species in 21 genera are recorded for Iran in this catalogue. After the catalogue one genus and six species have been added to the list and three new species have been described (Lelej & Osten, 2004). During a field survey in the western provinces of Iran including Ilam and Azarbaijan-e-Sharghi (=East Azarbaijan), eight species were collected by a light trap or directly by hand from the ground. One species is a new for the science and four species are newly recorded from Iran. All material has been identified by A. Lelej. Some additional material from the collection of the Zoological Institute (St. Petersburg) has been studied also. The distribution of species follows to Lelej (2002). New records from Iran are asterisked (*). The genera and species are given in alphabetical order.

LIST OF THE SPECIES

Genus *Dentilla* Lelej in Lelej et Kabakov, 1980

TYPE SPECIES: *Mutilla erronea* André, 1902 (male), by original designation (junior subjective synonym of *Mutilla curtiventris* André, 1901 (female) according to Pagliano & Strumia, 2007).

Dentilla dichroa (Sichel et Radoszkowski, 1869)

SPECIMENS EXAMINED. Ilam Prov., Shirvan-Chardavol Research Station, light trap, VI.2004, 4♂ (B. Gharali).

DISTRIBUTION. Turkmenistan, Afghanistan, Iran, Iraq, Palestine, Egypt (Sinai).

REMARK. This species was recorded from southern provinces including Hormozgan, Bushehr and Fars (Lelej & Osten, 2004) and new material shows wider distribution which extend throughout Zagros mountain range in the west of Iran.

Genus *Myrmilla* Wesmael, 1851

TYPE SPECIES: *Mutilla distincta* Lepeletier, 1845 (female), by subsequent designation (Ashmead, 1903) (junior subjective synonym of *Mutilla calva* Villers, 1789 (female), according to Morawitz, 1865).

Myrmilla (Eurygnathilla) emiliae (Dalla Torre, 1897)

SPECIMENS EXAMINED. Azarbaijan-e-Sharghi Prov., Marand, Peyam, 22.VIII 2003, 2 ♀ (H. Lotfalizadeh).

DISTRIBUTION. Armenia, Azerbaijan, Iraq, Iran, Turkey, Turkmenistan.

REMARK. This species was recorded from Golestan Prov., North Iran (Lelej & Osten, 2004) and new finding extends its distribution toward the north-west.

**Myrmilla (Myrmilla) etzchmiadzini* (Radoszkowski, 1885)

SPECIMENS EXAMINED. Azarbaijan-e-Sharghi Prov., Marand, Peyam, 22.VIII 2003, 1 ♀ (H. Lotfalizadeh).

DISTRIBUTION. Armenia, Azerbaijan, *Iran (Azarbaijan-e-Sharghi).

**Myrmilla (Pseudomutilla) skorikovi* Lelej, 1985

SPECIMENS EXAMINED. Azarbaijan-e-Sharghi Prov., Marand, Peyam, 22.VIII 2003, 1 ♀ (H. Lotfalizadeh).

DISTRIBUTION. Russia (Astrakhan, Dagestan), Azerbaijan, Georgia, *Iran (Azarbaijan-e-Sharghi).

Genus *Physetopoda* Schuster, 1949

TYPE SPECIES: *Physetopoda insularis* Schuster, 1949 (male), by original designation (junior subjective synonym of *Mutilla rubrocincta* Lucas, 1849 (male), according to Brothers, 1983. The latter designation of *Mutilla montana* Panzer, 1805 by Pagliano & Strumia (2007) is invalid.

Physetopoda portschinskii (Radoszkowski, 1888)

SPECIMENS EXAMINED. Ilam Prov., Shirvan-Chardavol Research Station, light trap, VI 2004, 3 ♂ (B. Gharali).

DISTRIBUTION. Russia (south-east of European part), Armenia, Azerbaijan, Iran (Hormozgan, Kerman), Kazakhstan, Turkmenistan (south-west).

REMARK. This species was recorded from Hormozgan and Kerman provinces (Lelej & Osten, 2004) and new material extend its distribution to the west of Iran.

Genus *Smicromyrme* Thomson, 1870

GENDER: Masculine (Article 30.2.4 of the Code)

TYPE SPECIES: *Mutilla rufipes* Fabricius, 1787 (female), by monotypy (attributed to Latreille, 1792).

**Smicromyrme (Eremotilla) mesasiaticus* Lelej, 1985

SPECIMENS EXAMINED. Ilam Prov., Shirvan-Chardavol Research Station, light trap, VI 2004, 1 ♂ (B. Gharali).

DISTRIBUTION: Turkmenistan, Uzbekistan, *Iran (Ilam).


Smicromyrme (Erimyrme) zagrosensis Lelej, sp. n.

SPECIMENS EXAMINED. Holotype – ♂, Iran: Ilam Prov., Shirvan-Chardavol Research Station, light trap, VI 2004 (B. Gharali), deposited in the collection of the Institute of Biology and Soil Science, Vladivostok.

DESCRIPTION. MALE. Body length 8.8 mm. Head, mesosoma, except metasternum and medial ventral part of mesopleuron, tegulae, legs and palps ferruginous-red; antennae ferruginous-red with darkish three apical flagellomeres; mandibles ferruginous-red with black apical denticles; palps brownish-red; legs ferruginous-red with brownish femorae; metasoma black with brownish-red sternum 1, tergum 7 and chestnut sterna 2 and 8; mid- and hind spurs whitish. Wings hyaline with brown veins, forewing (distally of cells) slightly infuscated, more darkened to the anterior margin. Body and legs clothed with subappressed short and scattered long erect pale pubescence; metasomal segments 2-6 with apical rare whitish fringe denser on tergum 2; felt lines on tergum 2 and sternum 2 golden. Terga 3-5 with pale pubescence which not forms a band.

Head width 0.8X thorax width including tegulae, rounded behind the eyes (in dorsal view), occipital carina visible above. Clypeus weakly concave, smooth and shiny medially, anterior border with two denticles and narrow emargination between them (Fig. 2). Scape distinctly bicarinate beneath. Ocelli medial size (Fig. 2), ratio postocellar distance : oculo-ocellar distance 1.6X; distance between posterior ocellus and posterior head margin almost equal to oculo-ocellar distance. Antennal segment 3 length 1.5X its width, 2.8X antennal segment 2 (pedicellus), and 0.85X antennal segment 4, the latter equal to antennal segment 5 (Fig. 5). Maxillar palps short, approximately 2X mandibular length. Antenno-ocular carina well developed, arcuated. Mandible tridentate with curved upper carina, third denticle distinctly tuberculate beneath (Fig. 3); large ventral tooth beneath near the base not widened apically (Fig. 4). Frons and vertex with coarse dense deep punctures.

Mesosoma length 1.6X maximal pronotal width, humeral angles rounded. Para-scutal carinae of scutum developed. Posterior coxae carinate inside. Tegulae not protruded beyond scuto-scutellar suture, shining, glabrous, with a few punctures inside. Propodeum reticulate with larger cells dorsally. Relation of *R*-abscissae is 1.9 : 1.2 : 1.7 : 2.7.



Figs 1-6. Males of *Smicromyrme*. 1-5) *S. zagrosensis* sp. n., holotype: 1) genitalia, lateral view, 2) head, anterior view, 3) mandible, dorsal view, 4) mandible, lateral view, 5) antennal segments 1-5; 6) *S. stepposus*, genitalia, lateral view. Scale bar 1 mm for fig. 2, 0.5 mm for figs 1, 3-6.

Metasomal segment 1 carinate beneath, its length 0.9X its maximal width; segment 2 with long lateral felt lines on tergum and very short ones on sternum; tergum 2 with dense large punctures. Tergum 7 weakly rounded apically, with large dense punctures, medially smooth and shiny. Genitalia laterally as Fig. 1.

FEMALE unknown.

DISTRIBUTION. Iran (Ilam).

DISCUSSION. A new species belongs to subgenus *Erimyrme* Lelej, 1985 with tridentate mandibles and concave clypeus in the males. In this subgenus a new species more related with *Smicromyrme kermanensis* (Lelej, 1984) and *S. stepposus* Lelej, 1984 by having the gonostyli with internal tuft of setae. A new species differs from *S. kermanensis* which has ferruginous-red mesosoma, legs and brownish-red head by having smaller ocelli (ratio *POD* : *OOD* 1.7X in *S. kermanensis*), by having medial smooth shiny area on metasomal tergum 7 (punctate throughout in *S.*

kermanensis), by having short felt line on metasomal sternum 2 (only a few small punctures in *S. kermanensis*), by having widened apically volsella (more or less parallelsided in *S. kermanensis*). *S. zagrosensis* sp. n. differs from *S. stepposus* by having ferruginous-red mesosoma (at least propodeum, mesopleuron ventrally, metapleuron, and legs black in *S. stepposus*), by having ferruginous-red head (black in *S. stepposus*), by having medial smooth shiny area on metasomal tergum 7 (punctate throughout in *S. stepposus*), by having another shape of volsella (Fig. 1 vs. Fig. 6).

ETYMOLOGY. The specific name is a Latin adjective derived from the Zagros Mountains, referring to the region where the species has been found.

Genus *Tropidotilla* Bischoff, 1920

TYPE SPECIES: *Mutilla littoralis* Petagna, 1787 (female), by original designation (misspelled as *littoralis*).

**Tropidotilla cypriadis* Invrea, 1940

SPECIMENS EXAMINED. Ilam Prov., Shirvan-Chardavol Research Station, light trap, VI 2004, 1♂ (B. Gharali).

DISTRIBUTION. Cyprus, Iraq, *Iran (Ilam).

ACKNOWLEDGEMENTS

We are grateful to V.M. Loktionov (Institute of Biology and Soil Science, Vladivostok, Russia) for the preparing of the figures, S.A. Belokobylskij (Zoological Institute, St. Petersburg, Russia) for the loan of material. The second author wishes to thank J. Hosseinzadeh for providing facilities during field trip.

The work was supported in part by the Russian Foundation for Basic Research (grant No 08-04-00184, A.S. Lelej principal investigator).

REFERENCES

- Brothers, D.J. 1983. Identity and classification of *Physetopoda*, *Chaetotilla* and *Paramyrmex*. – Journal of the Kansas Entomological Society 56(3): 441–445.
- Brothers, D.J. 1989. Alternative life-history styles of mutillid wasps (Insecta: Hymenoptera). – In: Bruton, M.N. (Ed.). Alternative life-history styles of animals. Dordrecht: Kluwer Academic Publishers. P. 279–291.
- Brothers, D.J. & Finnamore, A.T. 1993. Superfamily Vespoidea. – In: Goulet, H., Huber, J.T. (Eds): Hymenoptera of the World: An identification guide to families. Ottawa. P. 161–278, Figs 46–92.
- International Commission on Zoological Nomenclature (ICZN). 1999. International Code of Zoological Nomenclature. Fourth Edition. London: ITZN. i–xxx + 1–306.
- Lelej, A.S. 2002. Catalogue of the Mutillidae (Hymenoptera) of the Palaearctic Region. Vladivostok: Dalnauka, 172 p.
- Lelej, A.S. 2007. Biogeography of mutillid wasps (Hymenoptera, Mutillidae). – In: Rasnitsyn A.P. & Gokhman V.E. (Eds). Studies on Hymenopterous Insects. Collection of Scientific Papers. Moscow: KMK Scientific Press Ltd. P. 82–111. (In Russian).

Lelej, A.S., Brothers, D.J. 2008. The genus-group names of Mutillidae (Hymenoptera) and their type species, with a new genus, new name, new synonymies, new combinations and lectotypifications. – *Zootaxa* 1889: 1–79.

Lelej, A.S., T. Osten, 2004. To the knowledge of the mutillid and bradynobaenid wasps of Iran (Hymenoptera: Mutillidae, Bradynobaenidae). – *Proceeding of the Russian Entomological Society, St. Petersburg* 75(1); 253–262.

Morawitz, A. 1865. Über eine neue, oder vielmehr verkannte Form von Männchen unter den Mutillen, nebst einer Übersicht der in Europa beobachteten Arten. – *Bulletin de l'Académie Impériale des Sciences de Saint-Petersbourg*, 8: 82-141. (Reprinted in *Mélanges Biologiques tirés du Bulletin de l'Académie Impériale des Sciences de St.-Petersbourg*, 4, 671–756).

Pagliano, G. & Strumia, F. 2007. Contributo alla conoscenza dei Mutillidae italiani (Hymenoptera, Scolioidea). – *Bollettino del Musei Regionale di Scienze Naturali di Torino*, 24(1): 25–110.

SHORT COMMUNICATION

M. Yu. Proshchalykin. The bees of the family Melittidae (Hymenoptera, Apoidea) of Transbaikalia. – Far Eastern Entomologist. 2008. N 191: 8-9.

М. Ю. Прошалыкин. Пчелы семейства Melittidae (Hymenoptera, Apoidea) Забайкалья // Дальневосточный энтомолог. 2008. N 191. С. 8-9.

The melittid fauna of the world numbers 154 species in 14 genera. The family is rather small, greatest diversity is in the warm xeric areas [1]. Most species are oligolectic or at least have strong floral preferences, and some, *Macropis* and *Rediviva*, collect oils (to mix with pollen) [1]. There are nine species in three genera in the Russian Far East [2].

The family Melittidae was unknown in Transbaikalia. During an examination the material of the bees deposited in the collections of the Institute of Biology and Soil Sciences, Russian Academy of Sciences (Vladivostok) and Zoological Institute, Russian Academy of Sciences (St. Petersburg), four species in three genera were found and identified. All of them are newly recorded from this region. All distributed in Transbaikalia species are found in the Russian Far East. The distribution of melittids follows M. Proshchalykin [2].

The present investigation has been supported in part by the grants of the Far Eastern Branch of Russian Academy of Sciences No 06-III-B-195, No 06-III-A-06-138, No 06-III-A-06-147 and by the grant of the RFBR No 06-04-81008.

LIST OF THE SPECIES

***Macropis (Macropis) dimidiata amurensis* Popov, 1958**

MATERIAL. Russia: Buryatiya, Torei, 15.VII 1961, 1♂; Temnik river, 17.VII 1961, 2♀; Zabaikalskii krai, Tsasuchei, 16.VII 1990, 2♀, 2♂ (Romankova).

DISTRIBUTION. Russia: Buryatiya (new record), Zabaikalskii krai (former Chitinskaya oblast) (new record), Primorskii krai, Khabarovskii krai, Amurskaya oblast.

***Dasygoda altercator* (Harris, 1780)**

MATERIAL. Russia: Buryatiya, Ivolginskii datsan, 24.VII 1990, 3♂ (Romankova); Ulan-Ude, 26.VII 1990, 2♀, 1♂ (Romankova); Kalenovo, 28.VII 1990, 1♀ (Romankov); Gusinoe lake, Baraty, 26.VII 2007, 10♀, 20♂ (Lelej, Proshchalykin, Loktionov); Dzhida, 28.VII 2007, 2♀, 2♂ (Lelej, Proshchalykin, Loktionov); Naushki, Selenga river, 30.VII 2007, 11♀, 7♂ (Lelej, Proshchalykin, Loktionov).

DISTRIBUTION. Russia: Buryatiya (new record), Primorskii krai, Siberia, European part. – Mongolia, Central Asia, Caucasus, Europe, North Africa.

***Dasygoda japonica* Cockerell, 1911**

MATERIAL. Russia: Zabaikalskii krai, 20 km SSE Krasnokamenska, 3–5.VIII 2007, 11♂ (Lelej, Proshchalykin, Loktionov).

DISTRIBUTION. Russia: Zabaikalskii krai (new record), Primorskii krai, Khabarovskii krai. – Japan (Honshu, Kyushu), Korea, North-East China.

***Melitta tricincta* Kirby, 1802**

MATERIAL. Russia: Buryatiya, Peschanka, Troitskosavsk (=Kyakhta), 15.VIII–1.IX 1927, 7♂ (Mikhno); Ust-Kiran, Kiran river, 10.VII 1929, 1♂ (Mikhno); Ivolginskii datsan, 17.VI 1991, 2♂ (Romankova); Gusinoe lake, Baraty, 26.VII 2007, 1♀, 106♂ (Lelej, Proshchalykin, Loktionov); Dzhida, 28.VII 2007, 5♂ (Lelej, Proshchalykin, Loktionov); Naushki, Selenga river, 30.VII 2007, 1♂ (Lelej, Proshchalykin, Loktionov); Zabaikalskii krai, 20 km SSE Krasnokamensk, 3-5.VIII 2007, 18♂ (Lelej, Proshchalykin, Loktionov).

DISTRIBUTION. Russia: Buryatiya (new record), Zabaikalskii krai (new record), Khabarovskii krai, Yakutiya, European part. – Europe.

1. Michener, Ch.D. 2007. The Bees of the World. Baltimore, London: John Hopkins University Press. Second edition. 953 p.

2. Proshchalykin, M.Yu. 2007. Family Melittidae. – In: Lelej, A.S., Belokobylskij, S.A., Kasparyan, D.R., Kupianskaya, A.N. & Proshchalykin, M.Yu. (Eds). *Opredelitel' nasekomyh Dal'nego Vostoka Rossii* [Key to the insects of Russian Far East]. Vol. 4. Neuropteroidea, Mecoptera, Hymenoptera. Pt. 5. Vladivostok: Dal'nauka. P. 886-889. (In Russian).

Author's address:

Institute of Biology and Soil Science, FEB
RAS, Vladivostok-22, 690022, Russia

FAR EASTERN ENTOMOLOGIST
2008

CONTENTS

	N of issue	Pages	Date of issue
Yu. A. Tshistjakov. Far Eastern species of the genus <i>Achlya</i> Billberg, 1820 (Lepodoptera, Drepanidae: Thyatirinae) with notes on their synonymy	180	1-10	Jan.
V. G. Bezborodov, S. A. Shabalin and E. S. Koshkin. Distribution of <i>Anthracophora rusticola</i> Burmeister, 1842 (Coleoptera: Scarabaeidae, Cetoniinae) in Russia	180	11-12	Jan.
I. Trisnawati and K. Nakamura. Abundance, diversity and distribution of above-ground arthropods collected by window traps from satoyama in Kanazawa, Japan: an order level analysis	181	1-23	Feb.
Dahelmi, S. Salmah, I. Abbas, N. Fitriana, S. Nakano and K. Nakamura. Duration of immature stages of eleven swallowtail butterflies (Lepidoptera: Papilionidae) in West Sumatra, Indonesia	182	1-9	Mar.
A. L. Ozerov and M. Iwasa. A new species of the genus <i>Toxopoda</i> Macquart, 1851 (Diptera: Sepsidae) from India	182	10-11	Mar.
M. G. Krivosheina. A new genus and two new species of the shore-flies (Diptera: Ephydriidae) from Thailand	183	1-8	Mar.
A. L. Ozerov. Two new species of the genus <i>Norellia</i> Robineau-Desvoidy, 1830 (Diptera, Scathophagidae) from Altai	184	1-4	Apr.
Yu. A. Tshistjakov. A review of the subfamily Nolinae (Lepidoptera, Nolidae) of the Russian Far East	185	1-19	May
P. G. Nemkov. A key to the digger wasps of the genus <i>Mellinus</i> (Hymenoptera, Crabronidae, Mellininae) of the Russia	185	19-20	May
D. S. Aristov. Revision of the Permian genus <i>Parasylyviodes</i> Martynov, 1940 (Grylloblattida: Liomopteridae)	186	1-5	June
E. S. Koshkin. New records of the nolid-, noctuid-, and geometrid-moths (Lepidoptera: Nolidae, Noctuidae, Geometridae) from the Russian Far East	186	6-7	June
S. Yu. Storozhenko. <i>Uvarovina venosa</i> (Fischer-Waldheim, 1839) is newly recorded katydid species (Orthoptera, Tettigoniidae) from Buryatia	186	8	June
M. Yu. Proshchalykin. The bees of family Colletidae (Hymenoptera, Apoidea) of Transbaikalia	187	1-9	Aug.
P. G. Nemkov. New records of the digger wasps (Hymenoptera: Sphecidae, Crabronidae) from the Asian part of Russia	187	10-11	Aug.

A. V. Shavrin. Apateticinae Fauvel, 1895 is a new for the Russia subfamily of the rove beetles (Coleoptera, Staphylinidae)	187	12	Aug.
D. S. Aristov. New Grylloblattida (Insecta) from the Middle and Upper Permian of the Russia	188	1-7	Sep.
S. Yu. Storozhenko. <i>Diestrammena coreana</i> (Yamasaki, 1969) is newly recorded cave-cricket species (Orthoptera, Raphidophoridae) from Russia	188	8	Sep.
E.V. Mikhailjova. A new species of the millipede genus <i>Maritimosoma</i> (Diplopoda, Diplomaragnidae) from the Russian Far East	189	1-4	Oct.
A. S. Lelej and V. M. Loktionov. A review of the genus <i>Poecilagenia</i> Haupt, 1927 (Hymenoptera, Pompilidae) of the Russia with the world catalogue of the species	190	1-16	Nov.
A. S. Lelej, B. Gharali, H. Lotfalizadeh. New records of velvet ants (Hymenoptera: Mutillidae) from the west of Iran	191	1-7	Dec.
M. Yu. Proshchalykin. The bees of the family Melittidae (Hymenoptera, Apoidea) of Transbaikalia	191	8-9	Dec.

INSTRUCTIONS FOR AUTHORS

Far Eastern Entomologist is journal publishing original papers on entomology, including taxonomy, systematic, morphology, phylogeny, as well biology, ecology and biogeography. Reviews, comprehensive or revisionary studies of the insects thought other East Asia are especially welcome and will be given first priority for publication. Faunistic papers based on materials from the Russian Far East may be submitted also. Submission of a manuscript to Far Eastern Entomologist implies that the report is original, unpublished and is not being considered for publication elsewhere. Papers in languages other than English are not accepted.

Manuscripts must be type-written, double-spaced on one side of the standard-sized (A4, 21x31 sm.) white paper and submitted in one (text and figures) copy in a fully corrected form. Articles should be concise and the number of tables and figures limited to what is strictly necessary. Manuscripts should not exceed 16 pages (including figures and tables); additional printed pages are at the expense of the author(s).

Manuscripts should be prepared in accordance with the style and format of recent issues. (Current issues of Far Eastern Entomologist should be checked for style and format). An abstract should be followed by Key Words (2-7) and include no more than 100 words totally. Cite the author and year of publication of genera and species on first mention. The names of genera and species should be underlined. New description must conform with the current edition of the Code of Zoological Nomenclature. If a new taxon is described, the institution or museum where the type material is deposited must be indicated. The description of new taxa on types deposited in personal collection will not be accepted.

References in the text, as follows: "Bey-Bienko (1932) states..." or "Bey-Bienko (1932: 25) states..." when the author wishes to refer to a specific page, or "(Bey-Bienko, 1932)" as the author of a statement. Joint authors must be connected by "&" in both the text and the references. When there are more than two authors use "et al.," (Bey-Bienko et al., 1932) in the text. If journal names are not spelled out completely they should follow a consistent and accepted format.

Size of figures (including plates) as published should be less than 115 x 165mm. Illustration should be numbered in a single series throughout in Arabic numerals. Tables and legends must be typed on separate sheets and should be self-explanatory.

The following transliterations of Russian alphabet should be used:

A - a	E - e	K - k	П - p	Ф - f	Щ - shch
Б - b	Ж - zh	Л - l	Р - r	Х - kh	Ы - y
В - v	З - z	М - m	С - s	Ц - ts	Э - e
Г - g	И - i	Н - n	Т - t	Ч - ch	Ю - yu
Д - d	Й - i	О - o	У - u	Ш - sh	Я - ya

Manuscripts submitted are subject to review and editing by two anonymous reviewers and the Editor-in-Chief.

Twenty reprints of each article are provided free of charge to the first author. An order form and prices for additional reprints will be sent with the proofs.

Inquiries regarding content, subscription, manuscripts and copies should be sent to editor: S.Yu. Storozhenko, Institute of Biology and Soil Science, Vladivostok, 690022, Russia.

We recommend the electric submission of a complete manuscript to our office (entomol@ibss.dvo.ru).

© Far Eastern entomologist (Far East. entomol.) Journal published since October 1994.

Editor-in-Chief: S.Yu. Storozhenko

Editorial Board: A.S. Lelej, V.S. Sidorenko, N.V. Kurzenko, P.G. Nemkov

Address: Institute of Biology and Soil Science, Far East Branch of Russian Academy of Sciences, 690022, Vladivostok-22, Russia.

E-mail: entomol@ibss.dvo.ru

web-site: <http://www.biosoil.ru/fee>